

# Report Card Methods

## How are the grades calculated?

# **Freshwater Methods**

Freshwater monitoring is carried out twice a year, during spring (pre-wet season) and autumn (post wet season), at 135 representative sites across South East Queensland. Five ecological indicators are used to assess the health of freshwater ecosystems: physical and chemical, nutrient cycling, ecosystem processes, macroinvertebrates and fish.

#### Aquatic Macroinvertebrates

Aquatic macroinvertebrates (insects, crustaceans, snails, etc.) are common, widespread and easily sampled. They vary in sensitivity to disturbance and reflect environmental conditions, and thus stream health, over time. Sampling methods used are based on those used for the Queensland AusRivAS (Australian River Assessment System) program.

#### The three indices used are:

- Number of macroinvertebrate taxa
- PET richness (number of stonefly, mayfly and caddisfly families)
- SIGNAL (Stream Invertebrate Grade Number Average Level).

#### **Ecosystem Processes**

Measuring the rate of the production and decomposition of organic matter reflects the vigour or 'pulse' of a stream and indicates if it is healthy or unhealthy.

- The three indices used are:
- Ratio of <sup>13</sup>C to <sup>12</sup>C stable isotopes
- Respiration (R<sub>24</sub>)
- Gross Primary Production (GPP).

#### Nutrient Cycling

This describes the processing of nitrogen within a stream and the sensitivity of the stream to the input of nutrients.

#### The index used is:

Ratio of <sup>15</sup>N to <sup>14</sup>N stable isotopes.

### Fish

Fish communities reflect a range of environmental disturbances and provide a measure of stream condition due to their mobility, long life and position near the top of the food chain. Sampling of fish is carried out using a combination of electrofishing and seine netting.

The three indices used are:

- Percentage of Native Species Expected (PONSE)
- Ratio of Observed to Expected native species (O/E)
- Proportion of alien fish.

#### Physical and Chemical

Physical and chemical measures are important for monitoring direct changes in water quality and aiding in the interpretation of other measures of stream health.

The six indices used are:

- pH
- Conductivity
- Diel (24hr) range and maximum temperature
- Diel range and minimum dissolved oxygen.

## Freshwater Report Card Generation

- Results for each site are assessed against regional Ecosystem Health Guidelines for the corresponding stream type and standardised scores (ranging between 0 and 1) are derived.
- 2. The standardised scores for each of the indices within each indicator type are averaged to produce five summary scores per site.
- The indicator scores for all sites within a reporting area are averaged to produce five summary scores per reporting area.
- The values for the five indicator types are then averaged to give a single value for each reporting area.


Utrients

- The values for each reporting area are then averaged across seasons (spring and autumn).
- Catchments are then ranked based on these scores and Report Card grades are assigned.



# **Estuarine/Marine Methods**

The Estuarine and Marine Report Card Grade is calculated by combining an Ecosystem Health Index (EHI) and a Biological Health Rating (BHR) to produce a single value of ecosystem health.



rial should be addressed to Healthy Waterways.

Email: hwwinfo@healthywaterways.org

www.healthywaterways.org